

Exhibit No.: _____
Issues: Need for the Project
Witness: Donald W. Shaw
Sponsoring Party: Show-Me Concerned Land Owners
Type of Exhibit: Rebuttal Testimony
Case No.: EA-2016-0358
Date Testimony Prepared: January 24, 2017

**BEFORE THE PUBLIC SERVICE COMMISSION
OF THE STATE OF MISSOURI**

FILED
April 4, 2017
Data Center
Missouri Public
Service Commission

In the Matter of the Application of Grain Belt Express)
Clean Line LLC for a Certificate of Convenience and)
Necessity Authorizing it to Construct, Own, Operate,)
Control, Manage, and Maintain a High Voltage, Direct) Case No. EA-2016-0358
Current Transmission Line and an Associated Converter)
Station Providing an Interconnection on the Maywood-)
Montgomery 345 kV Transmission Line)

**REBUTTAL TESTIMONY OF
DONALD W. SHAW
ON BEHALF OF THE
SHOW ME CONCERNED LANDOWNERS
JANUARY 24, 2017**

Shaw Me Exhibit No. 402
Date 3-23-17 Reporter TS
File No. EA-2016-0358

1 **BEFORE THE PUBLIC SERVICE COMMISSION**
2 **OF THE STATE OF MISSOURI**
3 **REBUTTAL TESTIMONY OF**
4 **DONALD W. SHAW**
5 **ON BEHALF OF THE**
6 **SHOW ME CONCERNED LANDOWNERS**
7 **JANUARY 24, 2017**

8
9 **I INTRODUCTION**

10 **Q. Please state your name and address?**

11 A. My name is Donald W. Shaw. My address is 2409 Bowe Lane Dr., Jefferson City,
12 Missouri 65109.

13 **Q. On behalf of what party in this case are you testifying?**

14 A. I am testifying on behalf of the Show Me Concerned Landowners.

15 **Q. What is your education and professional background?**

16 A. I retired from Central Electric Power Cooperative. Prior to my retirement, I was Chief
17 Executive Officer of Central.

18 I attended the University of Missouri – Rolla and received a B.S. in Electrical
19 Engineering in 1969, a M.S. in Electrical Engineering in 1971, and a M.S. in Engineering
20 Management in 1978. My concentration was in Electrical Power Systems.

21 In 1971, I was employed by Missouri Public Service Company as a Substation
22 Design Engineer. In 1973, I took a position with Central Electric Power Cooperative
23 (“Central”), where I remained until I retired in 2015. I held several positions at Central,
24 including substation and high voltage transmission design positions and Manager of

1 Engineering and Technical Services. In the latter position, I was responsible for
2 transmission line right-of-way acquisition. When necessary, I represented Central as an
3 expert witness in eminent domain proceedings in circuit court. I served as CEO/General
4 Manager of Central from 1993-2015, and in that capacity, I also served as a member of
5 the Board of Directors of Associated Electric Cooperative (“AECI”), headquartered in
6 Springfield, Missouri. AECI owns the power generation facilities and controls the high
7 voltage transmission lines in the cooperative system. As an AECI director I attended
8 monthly board meetings and oversaw the planning, construction and operation of the
9 power plants and high voltage transmission lines. From 1993 through my retirement in
10 2015, I was ultimately responsible for all business decisions of Central. I was also
11 involved in all significant business decisions of AECI.

12 **II. NEED FOR THE PROJECT**

13 **Q. In your role as an expert witness in eminent domain proceedings, what was your**
14 **function?**

15 A. As Manager of the department that performed the technical studies for Central’s power
16 system improvements and made the final recommendation to the Central board to
17 approve the location of substation and transmission line additions, I was Central’s expert
18 witness that testified regarding the need for new facilities necessary for adequate and
19 reliable power system operation. This expert opinion was necessary for the Circuit Court
20 to find that proper technical studies by qualified engineers had been done and that the
21 rights being sought were needed to provide adequate and reliable electric service to
22 Missouri electrical consumers. My guiding principle was whether the new facilities were
23 necessary for the public benefit.

1 **Q. What preliminary steps did you undertake to determine whether a new**
2 **transmission line was needed?**

3 A. In anticipation of presenting a request to the Board of Directors, my department would
4 undertake studies to determine the need for additional facilities to providing basic electric
5 services as well as what property rights would be needed to support those facilities. My
6 department would look at many options for fulfilling the proposed “need” under
7 consideration. We did the study on a system wide basis, taking into consideration all
8 options for fulfilling the need for additional service. Oftentimes, the need could be
9 satisfied by simply using the facilities that already existed. One common approach was
10 to increase line operating voltage levels. In other words, some studies simply indicated
11 that there was no need for new facilities. If there was a need for the facilities, we would
12 present our studies to the Board of Directors for their approval.

13 We also used the studies in the Circuit Court proceedings, provided the Board of
14 Directors approved the new facilities. The evidence presented to the court always
15 included testimony regarding the public need for the proposed facility as well as the
16 evaluation of possible alternatives that were considered when reaching a conclusion
17 regarding the best overall solution.

18 **Q. Based on your business judgment and experience with Central and AECL, is there a**
19 **need for the Grain Belt Express project?**

20 A. No, there is not. While I have not done the engineering analysis of the project, it is clear
21 to me from the surrounding circumstances that there is no need for the project. There are
22 no load serving entities that have committed to purchasing service from the project. No

1 load serving entity has intervened in this case other than the Missouri Joint Municipal
2 Electric Utility Commission. I conclude that there has been no request for service
3 requiring the project, and, therefore, there is no need.

4 While certain wind generators have requested service from the GBX project, there
5 is no binding commitment to provide or pay for service. These requests for service are
6 merely a request for an opportunity to sell service. They are no more and no less than a
7 job seeker submitting his resume to a potential employer. Neither entity has an
8 obligation under such requests. The requests have no economic value. They are not
9 requests from entities which have an obligation to serve load.

10 In addition, neither MISO nor SPP have included the project as a transmission
11 facility in their study process. There, therefore, has been no system wide study to
12 determine the best way to respond to the “need” Clean Line claims. There has been no
13 assessment that the so called need can be handled by other system changes such as
14 increasing line operating voltage levels or system reconfigurations.

15 **Q. Does the contract between Clean Line and MJMEUC change your opinion?**

16 **A.** No, it does not. I agree with Glen Justis. There is no real economic value to MJMEUC
17 contract. Clean Line has no obligation to build the line. If they do, MJMEUC has no
18 obligation to take service from the GBX project. There is no commitment to provide or
19 take service from either party. As a cooperative executive, I would not have made a
20 business judgment based on the contract. I cannot see how either Clean Line or
21 MJMEUC can base any utility planning decision on the contract. The contract appears to
22 be designed as a means to acquire this Commission’s approval of the GBX project.

1 **Q. As an employee of Central, what education or training did you receive on climate**
2 **science?**

3 A. As an employee of Central Electric Power Cooperative and a board member of AECL, I
4 attended several seminars that included climate change presentations and discussions. I
5 had the opportunity to have one on one conversations with climate experts from various
6 Universities, including the Arizona State University and the University of Missouri. I
7 have also done independent research and made climate change presentations.

8 **Q. Do you agree with the common popular opinion that there is man made climate**
9 **change?**

10 A. No, I do not. First, let me be clear there is no consensus on man made climate change.
11 There does seem to be some consensus among government entities that there is
12 something to man made climate change. However, even that seems to be eroding. It is
13 simply not the case that man made climate change justifies a need for the GBX project. I
14 have attached supporting documents as schedules to my testimony with their supporting
15 links where possible. My first reference here is [Schedule DWS-1](#).

16 **Q. Please elaborate.**

17 A. There is no verifiable link between fossil fuel combustion and the accompanying release
18 of CO₂ into the atmosphere with climate change or global warming. Claims of global
19 warming and climate change are based on computer models. These models are based on
20 assumptions that have not yet been verified. As an example, according to computer
21 models, the earth should warm as CO₂ levels rise in the atmosphere. For the last 18
22 years, CO₂ levels have been rising, but satellite temperature measurements have recorded

1 no discernible temperature increases as the models have predicted. Legitimate science
2 requires that hypotheses be verified with conclusive data. Many of the projections made
3 by the computer models have been refuted or not supported by the data. My reference for
4 this is attached as [Schedule DWS-2](#).

5 **Q. If there were some measurable temperature change resulting from the emission of**
6 **CO2 into the atmosphere, what would the impact be in terms of climate change**
7 **from the GBX project?**

8 A. Allow me to start by quantifying the amount of CO₂ that exists in the atmosphere. The
9 air we breathe is about 78% Nitrogen 21% Oxygen and the remaining 1% is trace gasses,
10 including CO₂ which is about 0.04% (4 one hundredths of 1%). The amount of CO₂ in
11 our atmosphere is not even close to the CO₂ levels in our atmosphere at certain period in
12 our history. During certain periods in our history, CO₂ levels have been as much as 10
13 times higher than they are now. We have seen a small increase in CO₂ levels since the
14 year 1990 due to deforestation and increased use of fossil fuels. So it can be safely
15 concluded that CO₂ levels is not the overriding factor in our climate. Attached are two
16 temperature graphs over time as Schedule DWS-3.

17 In addition, China emits about 2 times the amount of CO₂ into the atmosphere that
18 the U.S. does. In 2014, worldwide CO₂ emissions totaled 35,699 million metric tons. The
19 United States was responsible for about 5,300 MMT and the US coal and natural gas
20 plants were responsible for about 2,000 MMT or 2,000/35,99 (5.5%) of the worldwide
21 total. Stated another way, if ALL of the US coal and natural gas generation was
22 eliminated worldwide CO₂ emissions would fall 5.5%.

1 Even if we could produce such a reduction in CO₂ emissions, the result would not
2 be good. The operation of the US transmission system has evolved with generation
3 sources and loads distributed in a manner that has operated in a stable and reliable
4 manner. A shift in the distribution of generation resources necessary to produce this
5 emission reduction would have a catastrophic impact on operation of the transmission
6 system. But it would have an immeasurably small impact on CO₂ emissions and virtually
7 no impact on the earth's temperature which has been basically stable for the past 18
8 years. And certainly, the impact of the GBE project would be miniscule.

9 **Q. What are your conclusions regarding the GBE project?**

10 A. The Commission should deny the application for a CCN. There is no need for the project
11 to serve load within Missouri. There is no climate justification significant enough to
12 compromise the property rights of the landowners in the state of Missouri.

13 **Q. Does this conclude your testimony?**

14 A. Yes, it does.

**BEFORE THE PUBLIC SERVICE COMMISSION
OF THE STATE OF MISSOURI**


In the Matter of the Application of Grain Belt Express)
Clean Line LLC for a Certificate of Convenience and)
Necessity Authorizing it to Construct, Own, Control,) Case No. EA-2016-0358
Manage, Operate and Maintain a High Voltage, Direct)
Current Transmission Line and an Associated Converter)
Station Providing an Interconnection on the Maywood-)
Montgomery 345 kV Transmission Line)

AFFIDAVIT OF DONALD W. SHAW

STATE OF MISSOURI)
) ss
COUNTY OF Cole)

Donald W. Shaw, being first duly sworn on his oath, states:


1. My name is Donald W. Shaw.
2. Attached hereto and made a part hereof for all purposes is my Rebuttal Testimony on behalf of Show Me Concerned Landowners, consisting of 7 pages, having been prepared in written form for introduction into evidence in the above-captioned docket.
3. I have knowledge of the matters set forth therein. I hereby swear and affirm that my answers contained in the attached testimony to the questions therein propounded, including any attachments thereto, are true and accurate to the best of my knowledge, information and belief.


Donald W. Shaw

Subscribed and sworn before me this 23 day of January, 2017.

JOSHUA R. HANLIN
Notary Public - Notary Seal
STATE OF MISSOURI
County of Cole
My Commission Expires 9/15/2019
Commission # 15638231


Notary Public



Global Warming Petition

Home
Summary of Peer-Reviewed Research
Letter From Frederick Seitz
List of Signers By State
List of Signers By Name
Purpose of Petition
How Petition is Circulated
Instructions for Signing Petition
Qualifications of Signers

Summary of Peer-Reviewed Research

Most scientists have a detailed knowledge of their own narrow field of specialization, a general knowledge of fundamental science, an understanding of the scientific method, and a mental model that encompasses a broad range of scientific disciplines. This model serves as the basis of their thoughts about scientific questions.

When a scientist desires to refine his understanding of a specific scientific subject, he often begins by reading one or more review articles about that topic. As he reads, he compares the facts given in the review with his mental model of the subject, refining his model and updating it with current information. Review articles do not present new discoveries. The essential facts given in the review must be referenced to the peer-reviewed scientific research literature, so that the reader can check the assertions and conclusions of the article and obtain more detailed information about aspects that interest him.

A 12-page review article about the human-caused global warming hypothesis

Natale A. Marini
 David A. Oliver
 William B. Hardin
 Thomas O. Livingston
 Arthur J. Sherman
 M. Paul Reiter
 Vincent P. Rocco
 Ingrid Anderson
 Jerry A. Bradshaw
 Rudolph Neal Band
 Bettina Heinz
 George C. Pfaff Jr.
 Michael A. Crabb
 Arthur F. Widtfeldt
 Daniel Tao
 Theodore Lynn Rebstock
 Daniel Weiss
 Louis A. Williams, Jr
 David Flowers
 D. E. Maguire
 Donald R. Keys
 Vern S. Strubeck
 John P. Mihlbauer
 Loren Elwood Wiesner
 Don R. Morton
 Paris D. Svoronos
 Kent M. Mangold
 Michael A. Burke
 L. Gerald Marshall
 Theodore A. Ruppert
 Francis J. Nash Jr.
 Gary C. Prechter
 Gregory J. Brunetta
 Kenneth Lagrand
 Larry J. Pemberton
 Donald Jones Channin
 Thomas R. Stauffer
 Juris Vagners
 Michael S. Henson
 Donald J. Just
 David Thurmond Clark
 Bill J. Wright
 Ernest J. Andberg
 Dennis Skala
 C. Elaine Lane
 E. Brett Schafer
 Frederick H. Suydam
 Alan E. Munter
 R. S. Bennett
 Shawn B. Kendall
 Gustav Stolz Jr.
 Saul Kay
 Harendra Sakarlal Gandhi
 Dean Earl McFeron
 Frank J. Taverna
 Emmet Jones
 Ronald H. Duckstein Jr.
 Charles R. Galloway
 James G. McGee
 Mike Lauriente
 Lee J. Richard
 C. A. Brown
 Stuart Havenstrite
 David A. Hoecke
 Lowell C. Hanson
 Thomas Alexander Gleeson
 Tim F. Fridav

Schedule DWS-1

Frequently Asked Questions

is circulated with the petition. To view the entire article in [html](#), [150-dpi PDF](#), [300-dpi PDF](#), [600-dpi PDF](#), [Spanish](#) or figures alone in [powerpoint](#) or [flash](#), click on the appropriate item in this sentence.

Environmental Effects of Increased Atmospheric Carbon Dioxide

ARTHUR B. ROBINSON, NOAH E. ROBINSON, AND WILLIE SOON

Oregon Institute of Science and Medicine, 2251 Dick George Road, Cave Junction, Oregon 97523 [ar@oisom.org]

ABSTRACT A review of the research literature concerning the environmental consequences of increased levels of atmospheric carbon dioxide leads to the conclusion that increases during the 20th and early 21st centuries have produced no deleterious effects upon Earth's weather and climate. Increased carbon dioxide has, however, markedly increased plant growth. Predictions of harmful climatic effects due to future increases in hydrocarbon use and minor greenhouse gases like CO₂ do not conform to current experimental knowledge. The environmental effects of rapid expansion of the nuclear and hydrocarbon energy industries are discussed.

SUMMARY

Political leaders gathered in Kyoto, Japan, in December 1997 to consider a world treaty restricting human production of "greenhouse gases," chiefly carbon dioxide (CO₂). They feared that CO₂ would result in "human-caused global warming" - hypothetical severe increases in Earth's temperatures, with disastrous environmental consequences. During the past 10 years, many political efforts have been made to force worldwide agreement to the Kyoto treaty.

When we reviewed this subject in 1998 (1,2), existing satellite records were short and were centered on a period of changing interdecadal temperature trends. Additional experimental data have now been obtained, so better answers to the questions raised by the hypothesis of "human-caused global warming" are now available.




Figure 1: Surface temperatures in the Sargasso Sea, a 2 million square mile region of the Atlantic Ocean, with time resolution of 50 to 100 years and ending in 1975, as determined by isotope ratios of marine organisms retained in sediment at the bottom of the sea (3). The horizontal line is the average temperature for this 3,000-year period. The Little Ice Age and Medieval Climate Optimum were naturally occurring, extended intervals of climate departures from the mean. A value of 0.25 °C, which is the change in Sargasso Sea temperature between 1975 and 2006, has been added to the 1975 data in order to provide a 2006 temperature value.

The average temperature of the Earth has varied within a range of about 3°C during the past 3,000 years. It is currently increasing as the Earth recovers from a period that is known as the Little Ice Age, as shown in Figure 1. George Washington and his army were at Valley Forge during the coldest era in 1,500 years, but even then the temperature was only about 1° Centigrade below the 3,000-year average. The most recent part of this warming period is reflected by short-

Journal of American Physicians and Surgeons (2007) 12, 79-90.




Figure 2: Average length of 160 glaciers from 1700 to 2000 (4). The principal source of melt energy is solar radiation. Variations in glacier mass and length are primarily due to temperature and precipitation (5,6). This melting trend lags the temperature increase by about 20 years, so it produces the 6-fold increase in hydrocarbon use (7) even more than shown in the figure. Hydrocarbon use could not have caused this shortening trend.

ening of world glaciers, as shown in Figure 2. Glaciers regularly lengthen and shorten in delayed correlation with cooling and warming trends. Shortening lags temperature by about 20 years, so the current warming trend began in about 1800.

Atmospheric temperature is regulated by the sun, which fluctuates in activity as shown in Figure 3; by the greenhouse effect, largely caused by atmospheric water vapor (H₂O); and by other phenomena that are more poorly understood. While major greenhouse gas H₂O substantially warms the Earth, minor greenhouse gases such as CO₂




Figure 3: Arctic surface air temperature compared with total solar irradiance as measured by sunspot cycle amplitude, sunspot cycle length, solar equatorial rotation rate, fraction of penumbra spots, and decay rate of the 11-year sunspot cycle (8,9). Solar irradiance correlates well with Arctic temperature, while hydrocarbon use (7) does not correlate.

The factual information cited in this article is referenced to the underlying research literature, in this case by 132 references listed at the end of the article. Although written primarily for scientists, most of this article

can be understood without formal scientific training. This article was submitted to many scientists for comments and suggestions before it was finalized and submitted for publication. It then underwent ordinary peer review by the publishing journal.

The United Nations IPCC also publishes a research review in the form of a voluminous, occasionally-updated report on the subject of climate change, which the United Nations asserts is “authored” by approximately 600 scientists. These “authors” are not, however – as is ordinarily the custom in science – permitted power of approval the published review of which they are putative authors. They are permitted to comment on the draft text, but the final text neither conforms to nor includes many of their comments. The final text conforms instead to the United Nations objective of building support for world taxation and rationing of industrially-useful energy.

Environmental Effects of Increased Atmospheric Carbon Dioxide

ARTHUR B. ROBINSON, NOAH E. ROBINSON, AND WILLIE SOON

Oregon Institute of Science and Medicine, 2251 Dick George Road, Cave Junction, Oregon 97523 [atr@oism.org]

ABSTRACT A review of the research literature concerning the environmental consequences of increased levels of atmospheric carbon dioxide leads to the conclusion that increases during the 20th and early 21st centuries have produced no deleterious effects upon Earth's weather and climate. Increased carbon dioxide has, however, markedly increased plant growth. Predictions of harmful climatic effects due to future increases in hydrocarbon use and minor greenhouse gases like CO₂ do not conform to current experimental knowledge. The environmental effects of rapid expansion of the nuclear and hydrocarbon energy industries are discussed.

SUMMARY

Political leaders gathered in Kyoto, Japan, in December 1997 to consider a world treaty restricting human production of "greenhouse gases," chiefly carbon dioxide (CO₂). They feared that CO₂ would result in "human-caused global warming" – hypothetical severe increases in Earth's temperatures, with disastrous environmental consequences. During the past 10 years, many political efforts have been made to force worldwide agreement to the Kyoto treaty.

When we reviewed this subject in 1998 (1,2), existing satellite records were short and were centered on a period of changing intermediate temperature trends. Additional experimental data have now been obtained, so better answers to the questions raised by the hypothesis of "human-caused global warming" are now available.




Figure 1: Surface temperatures in the Sargasso Sea, a 2 million square mile region of the Atlantic Ocean, with time resolution of 50 to 100 years and ending in 1975, as determined by isotope ratios of marine organism remains in sediment at the bottom of the sea (3). The horizontal line is the average temperature for this 3,000-year period. The Little Ice Age and Medieval Climate Optimum were naturally occurring, extended intervals of climate departures from the mean. A value of 0.25 °C, which is the change in Sargasso Sea temperature between 1975 and 2006, has been added to the 1975 data in order to provide a 2006 temperature value.

The average temperature of the Earth has varied within a range of about 3°C during the past 3,000 years. It is currently increasing as the Earth recovers from a period that is known as the Little Ice Age, as shown in Figure 1. George Washington and his army were at Valley Forge during the coldest era in 1,500 years, but even then the temperature was only about 1° Centigrade below the 3,000-year average.

The most recent part of this warming period is reflected by short-




Figure 2: Average length of 169 glaciers from 1700 to 2000 (4). The principal source of melt energy is solar radiation. Variations in glacier mass and length are primarily due to temperature and precipitation (5,6). This melting trend lags the temperature increase by about 20 years, so it predates the 6-fold increase in hydrocarbon use (7) even more than shown in the figure. Hydrocarbon use could not have caused this shortening trend.

ening of world glaciers, as shown in Figure 2. Glaciers regularly lengthen and shorten in delayed correlation with cooling and warming trends. Shortening lags temperature by about 20 years, so the current warming trend began in about 1800.

Atmospheric temperature is regulated by the sun, which fluctuates in activity as shown in Figure 3; by the greenhouse effect, largely caused by atmospheric water vapor (H₂O); and by other phenomena that are more poorly understood. While major greenhouse gas H₂O substantially warms the Earth, minor greenhouse gases such as CO₂




Figure 3: Arctic surface air temperature compared with total solar irradiance as measured by sunspot cycle amplitude, sunspot cycle length, solar equatorial rotation rate, fraction of penumbral spots, and decay rate of the 11-year sunspot cycle (8,9). Solar irradiance correlates well with Arctic temperature, while hydrocarbon use (7) does not correlate.




Figure 4: Annual mean surface temperatures in the contiguous United States between 1880 and 2006 (10). The slope of the least-squares trend line for this 127-year record is 0.5 °C per century.

have little effect, as shown in Figures 2 and 3. The 6-fold increase in hydrocarbon use since 1940 has had no noticeable effect on atmospheric temperature or on the trend in glacier length.

While Figure 1 is illustrative of most geographical locations, there is great variability of temperature records with location and regional climate. Comprehensive surveys of published temperature records confirm the principal features of Figure 1, including the fact that the current Earth temperature is approximately 1 °C lower than that during the Medieval Climate Optimum 1,000 years ago (11,12).

Surface temperatures in the United States during the past century reflect this natural warming trend and its correlation with solar activity, as shown in Figures 4 and 5. Compiled U.S. surface temperatures have increased about 0.5 °C per century, which is consistent with other historical values of 0.4 to 0.5 °C per century during the recovery from the Little Ice Age (13-17). This temperature change is slight as compared with other natural variations, as shown in Figure 6. Three intermediate trends are evident, including the decreasing trend used to justify fears of "global cooling" in the 1970s.

Between 1900 and 2000, on absolute scales of solar irradiance and degrees Kelvin, solar activity increased 0.19%, while a 0.5 °C temperature change is 0.21%. This is in good agreement with estimates that Earth's temperature would be reduced by 0.6 °C through particulate blocking of the sun by 0.2% (18).

Solar activity and U.S. surface temperature are closely correlated, as shown in Figure 5, but U.S. surface temperature and world hydrocarbon use are not correlated, as shown in Figure 13.

The U.S. temperature trend is so slight that, were the temperature




Figure 5: U.S. surface temperature from Figure 4 as compared with total solar irradiance (19) from Figure 3.




Figure 6: Comparison between the current U.S. temperature change per century, the 3,000-year temperature range in Figure 1, seasonal and diurnal range in Oregon, and seasonal and diurnal range throughout the Earth.

change which has taken place during the 20th and 21st centuries to occur in an ordinary room, most of the people in the room would be unaware of it.

During the current period of recovery from the Little Ice Age, the U.S. climate has improved somewhat, with more rainfall, fewer tornados, and no increase in hurricane activity, as illustrated in Figures 7 to 10. Sea level has trended upward for the past 150 years at a rate of 7 inches per century, with 3 intermediate uptrends and 2 periods of no increase as shown in Figure 11. These features are confirmed by the glacier record as shown in Figure 12. If this trend continues as




Figure 7: Annual precipitation in the contiguous 48 United States between 1895 and 2006. U.S. National Climatic Data Center, U.S. Department of Commerce 2006 Climate Review (20). The trend shows an increase in rainfall of 1.8 inches per century – approximately 6% per century.




Figure 8: Annual number of strong-to-violent category F3 to F5 tornados during the March-to-August tornado season in the U.S. between 1950 and 2006. U.S. National Climatic Data Center, U.S. Department of Commerce 2006 Climate Review (20). During this period, world hydrocarbon use increased 6-fold, while violent tornado frequency decreased by 43%.




Figure 9: Annual number of Atlantic hurricanes that made landfall between 1900 and 2006 (21). Line is drawn at mean value.

did that prior to the Medieval Climate Optimum, sea level would be expected to rise about 1 foot during the next 200 years.

As shown in Figures 2, 11, and 12, the trends in glacier shortening and sea level rise began a century *before* the 60-year 6-fold increase in hydrocarbon use, and have not changed during that increase. Hydrocarbon use could not have caused these trends.

During the past 50 years, atmospheric CO₂ has increased by 22%. Much of that CO₂ increase is attributable to the 6-fold increase in human use of hydrocarbon energy. Figures 2, 3, 11, 12, and 13 show, however, that human use of hydrocarbons has not caused the observed increases in temperature.

The increase in atmospheric carbon dioxide has, however, had a substantial environmental effect. Atmospheric CO₂ fertilizes plants. Higher CO₂ enables plants to grow faster and larger and to live in drier climates. Plants provide food for animals, which are thereby also enhanced. The extent and diversity of plant and animal life have both increased substantially during the past half-century. Increased temperature has also mildly stimulated plant growth.

Does a catastrophic amplification of these trends with damaging climatological consequences lie ahead? There are no experimental data that suggest this. There is also no experimentally validated theoretical evidence of such an amplification.

Predictions of catastrophic global warming are based on computer climate modeling, a branch of science still in its infancy. The empirical evidence – actual measurements of Earth's temperature and climate – shows no man-made warming trend. Indeed, during four of the seven decades since 1940 when average CO₂ levels steadily increased, U.S. average temperatures were actually decreasing.




Figure 10: Annual number of violent hurricanes and maximum attained wind speed during those hurricanes in the Atlantic Ocean between 1944 and 2006 (22,23). There is no upward trend in either of these records. During this period, world hydrocarbon use increased 6-fold. Lines are mean values.




Figure 11: Global sea level measured by surface gauges between 1807 and 2002 (24) and by satellite between 1993 and 2006 (25). Satellite measurements are shown in gray and agree with tide gauge measurements. The overall trend is an increase of 7 inches per century. Intermediate trends are 9, 0, 12, 0, and 12 inches per century, respectively. This trend lags the temperature increase, so it predates the increase in hydrocarbon use even more than is shown. It is unaffected by the very large increase in hydrocarbon use.

While CO₂ levels have increased substantially and are expected to continue doing so and humans have been responsible for part of this increase, the effect on the environment has been benign.

There is, however, one very dangerous possibility.

Our industrial and technological civilization depends upon abundant, low-cost energy. This civilization has already brought unprecedented prosperity to the people of the more developed nations. Billions of people in the less developed nations are now lifting themselves from poverty by adopting this technology.

Hydrocarbons are essential sources of energy to sustain and extend prosperity. This is especially true of the developing nations, where available capital and technology are insufficient to meet rapidly increasing energy needs without extensive use of hydrocarbon fuels. If, through misunderstanding of the underlying science and through misguided public fear and hysteria, mankind significantly rations and restricts the use of hydrocarbons, the worldwide increase in prosperity will stop. The result would be vast human suffering and the loss of hundreds of millions of human lives. Moreover, the prosperity of those in the developed countries would be greatly reduced.

Mild ordinary natural increases in the Earth's temperature have occurred during the past two to three centuries. These have resulted in some improvements in overall climate and also some changes in




Figure 12: Glacier shortening (4) and sea level rise (24,25). Gray area designates estimated range of error in the sea level record. These measurements lag air temperature increases by about 20 years. So, the trends began more than a century before increases in hydrocarbon use.

the landscape, such as a reduction in glacier lengths and increased vegetation in colder areas. Far greater changes have occurred during the time that all current species of animals and plants have been on the Earth. The relative population sizes of the species and their geographical distributions vary as they adapt to changing conditions.

The temperature of the Earth is continuing its process of fluctuation in correlation with variations in natural phenomena. Mankind, meanwhile, is moving some of the carbon in coal, oil, and natural gas from below ground to the atmosphere and surface, where it is available for conversion into living things. We are living in an increasingly lush environment of plants and animals as a result. This is an unexpected and wonderful gift from the Industrial Revolution.

ATMOSPHERIC AND SURFACE TEMPERATURES

Atmospheric and surface temperatures have been recovering from an unusually cold period. During the time between 200 and 500 years ago, the Earth was experiencing the "Little Ice Age." It had descended into this relatively cool period from a warm interval about 1,000 years ago known as the "Medieval Climate Optimum." This is shown in Figure 1 for the Sargasso Sea.

During the Medieval Climate Optimum, temperatures were warm enough to allow the colonization of Greenland. These colonies were abandoned after the onset of colder temperatures. For the past 200 to 300 years, Earth temperatures have been gradually recovering (26). Sargasso Sea temperatures are now approximately equal to the average for the previous 3,000 years.

The historical record does not contain any report of "global warming" catastrophes, even though temperatures have been higher than they are now during much of the last three millennia.

The 3,000-year range of temperatures in the Sargasso Sea is typical of most places. Temperature records vary widely with geographical location as a result of climatological characteristics unique to those specific regions, so an "average" Earth temperature is less meaningful than individual records (27). So called "global" or "hemispheric" averages contain errors created by averaging systematically different aspects of unique geographical regions and by inclusion of regions where temperature records are unreliable.

Three key features of the temperature record – the Medieval Climate Optimum, the Little Ice Age, and the Not-Unusual-Temperature of the 20th century – have been verified by a review of local temperature and temperature-correlated records throughout the world (11), as summarized in Table 1. Each record was scored with respect to those queries to which it applied. The experimental and historical literature definitively confirms the primary features of Figure 1.

Most geographical locations experienced both the Medieval Climate Optimum and the Little Ice Age – and most locations did not

Table 1: Query	Yes	No	Yes/No	Two-Tailed Probability
Warm Climatic Anomaly 800-1300 A.D.?	88	2	7	> 99.99
Cold Climatic Anomaly 1300-1900 A.D.?	105	2	2	> 99.99
20th Century Warmest in Individual Record?	7	64	14	< 0.0001

Table 1: Comprehensive review of all instances in which temperature or temperature-correlated records from localities throughout the world permit answers to queries concerning the existence of the Medieval Climate Optimum, the Little Ice Age, and an unusually warm anomaly in the 20th century (11). The compiled and tabulated answers confirm the three principal features of the Sargasso Sea record shown in Figure 1. The probability that the answer to the query in column 1 is "yes" is given in column 5.




Figure 13: Seven independent records – solar activity (9); Northern Hemisphere, (13), Arctic (28), global (10), and U.S. (10) annual surface air temperatures; sea level (24,25); and glacier length (4) – all qualitatively confirm each other by exhibiting three intermediate trends – warmer, cooler, and warmer. Sea level and glacier length are shown minus 20 years, correcting for their 20-year lag of atmospheric temperature. Solar activity, Northern Hemisphere temperature, and glacier lengths show a low in about 1800.

Hydrocarbon use (7) is uncorrelated with temperature. Temperature rose for a century before significant hydrocarbon use. Temperature rose between 1910 and 1940, while hydrocarbon use was almost unchanged. Temperature then fell between 1940 and 1972, while hydrocarbon use rose by 330%. Also, the 150 to 200-year slopes of the sea level and glacier trends were unchanged by the very large increase in hydrocarbon use after 1940.

experience temperatures that were unusually warm during the 20th century. A review of 23 quantitative records has demonstrated that mean and median world temperatures in 2006 were, on average, approximately 1 °C or 2 °F cooler than in the Medieval Period (12).

World glacier length (4) and world sea level (24,25) measurements provide records of the recent cycle of recovery. Warmer temperatures diminish glaciers and cause sea level to rise because of decreased ocean water density and other factors.

These measurements show that the trend of 7 inches per century increase in sea level and the shortening trend in average glacier length both began a century before 1940, yet 84% of total human annual hydrocarbon use occurred only after 1940. Moreover, neither of these trends has accelerated during the period between 1940 and 2007, while hydrocarbon use increased 6-fold. Sea level and glacier records are offset by about 20 years because of the delay between temperature rise and glacier and sea level change.

If the natural trend in sea level increase continues for another two centuries as did the temperature rise in the Sargasso Sea as the Earth entered the Medieval Warm Period, sea level would be expected to rise about 1 foot between the years 2000 and 2200. Both the sea level and glacier trends – and the temperature trend that they reflect – are




Figure 14: Satellite microwave sounding unit (blue) measurements of tropospheric temperatures in the Northern Hemisphere between 0 and 82.5 N, Southern Hemisphere between 0 and 82.5 S, tropics between 20S and 20N, and the globe between 82.5N and 82.5S between 1979 and 2007 (29), and radiosonde balloon (red) measurements in the tropics (29). The balloon measurements confirm the satellite technique (29-31). The warming anomaly in 1997-1998 (gray) was caused by El Niño, which, like the overall trends, is unrelated to CO₂ (32).

unrelated to hydrocarbon use. A further doubling of world hydrocarbon use would not change these trends.

Figure 12 shows the close correlation between the sea level and glacier records, which further validates both records and the duration and character of the temperature change that gave rise to them.

Figure 4 shows the annual temperature in the United States during the past 127 years. This record has an upward trend of 0.5 °C per century. Global and Northern Hemisphere surface temperature records shown in Figure 13 trend upward at 0.6 °C per century. These records are, however, biased toward higher temperatures in several ways. For example, they preferentially use data near populated areas (33), where heat island effects are prevalent, as illustrated in Figure 15. A trend of 0.5 °C per century is more representative (13-17).

The U.S. temperature record has two intermediate uptrends of comparable magnitude, one occurring before the 6-fold increase in hydrocarbon use and one during it. Between these two is an intermediate temperature downtrend, which led in the 1970s to fears of an impending new ice age. This decrease in temperature occurred during a period in which hydrocarbon use increased 3-fold.

Seven independent records – solar irradiance; Arctic, Northern Hemisphere, global, and U.S. annual average surface air temperatures; sea level; and glacier length – all exhibit these three intermediate trends, as shown in Figure 13. These trends confirm one another. Solar irradiance correlates with them. Hydrocarbon use does not.

The intermediate uptrend in temperature between 1980 and 2006 shown in Figure 13 is similar to that shown in Figure 14 for balloon and satellite tropospheric measurements. This trend is more pronounced in the Northern Hemisphere than in the Southern. Contrary to the CO₂ warming climate models, however, tropospheric temperatures are not rising faster than surface temperatures.

Figure 6 illustrates the magnitudes of these temperature changes by comparing the 0.5 °C per century temperature change as the Earth recovers from the Little Ice Age, the range of 50-year averaged Atlantic ocean surface temperatures in the Sargasso Sea over the past 3,000 years, the range of day-night and seasonal variation on average

in Oregon, and the range of day-night and seasonal variation over the whole Earth. The two-century-long temperature change is small.

Tropospheric temperatures measured by satellite give comprehensive geographic coverage. Even the satellite measurements, however, contain short and medium-term fluctuations greater than the slight warming trends calculated from them. The calculated trends vary significantly as a function of the most recent fluctuations and the lengths of the data sets, which are short.

Figure 3 shows the latter part of the period of warming from the Little Ice Age in greater detail by means of Arctic air temperature as compared with solar irradiance, as does Figure 5 for U.S. surface temperature. There is a close correlation between solar activity and temperature and none between hydrocarbon use and temperature. Several other studies over a wide variety of time intervals have found similar correlations between climate and solar activity (15, 34-39).

Figure 3 also illustrates the uncertainties introduced by limited time records. If the Arctic air temperature data before 1920 were not available, essentially no uptrend would be observed.

This observed variation in solar activity is typical of stars close in size and age to the sun (40). The current warming trends on Mars (41), Jupiter (42), Neptune (43,44), Neptune's moon Triton (45), and Pluto (46-48) may result, in part, from similar relations to the sun and its activity – like those that are warming the Earth.

Hydrocarbon use and atmospheric CO₂ do not correlate with the observed temperatures. Solar activity correlates quite well. Correlation does not prove causality, but non-correlation proves non-causality. Human hydrocarbon use is not measurably warming the earth. Moreover, there is a robust theoretical and empirical model for solar warming and cooling of the Earth (8,19,49,50). The experimental data do not prove that solar activity is the only phenomenon responsible for substantial Earth temperature fluctuations, but they do show that human hydrocarbon use is not among those phenomena.

The overall experimental record is self-consistent. The Earth has been warming as it recovers from the Little Ice Age at an average rate of about 0.5 °C per century. Fluctuations within this temperature trend include periods of more rapid increase and also periods of temperature decrease. These fluctuations correlate well with concomitant fluctuations in the activity of the sun. Neither the trends nor the fluctuations within the trends correlate with hydrocarbon use. Sea level and glacier length reveal three intermediate uptrends and two downtrends since 1800, as does solar activity. These trends are climatically benign and result from natural processes.




Figure 15: Surface temperature trends for 1940 to 1996 from 107 measuring stations in 49 California counties (51,52). The trends were combined for counties of similar population and plotted with the standard errors of their means. The six measuring stations in Los Angeles County were used to calculate the standard error of that county, which is plotted at a population of 8.9 million. The “urban heat island effect” on surface measurements is evident. The straight line is a least-squares fit to the closed circles. The points marked “X” are the six unadjusted station records selected by NASA GISS (53-55) for use in their estimate of global surface temperatures. Such selections make NASA GISS temperatures too high.

ATMOSPHERIC CARBON DIOXIDE

The concentration of CO₂ in Earth's atmosphere has increased during the past century, as shown in Figure 17. The magnitude of this atmospheric increase is currently about 4 gigatons (Gt C) of carbon per year. Total human industrial CO₂ production, primarily from use of coal, oil, and natural gas and the production of cement, is currently about 8 Gt C per year (7,56,57). Humans also exhale about 0.6 Gt C per year, which has been sequestered by plants from atmospheric CO₂. Office air concentrations often exceed 1,000 ppm CO₂.

To put these figures in perspective, it is estimated that the atmosphere contains 780 Gt C; the surface ocean contains 1,000 Gt C; vegetation, soils, and detritus contain 2,000 Gt C; and the intermediate and deep oceans contain 38,000 Gt C, as CO₂ or CO₂ hydration products. Each year, the surface ocean and atmosphere exchange an estimated 90 Gt C; vegetation and the atmosphere, 100 Gt C; marine biota and the surface ocean, 50 Gt C; and the surface ocean and the intermediate and deep oceans, 40 Gt C (56,57).

So great are the magnitudes of these reservoirs, the rates of exchange between them, and the uncertainties of these estimated numbers that the sources of the recent rise in atmospheric CO₂ have not been determined with certainty (58,59). Atmospheric concentrations of CO₂ are reported to have varied widely over geological time, with peaks, according to some estimates, some 20-fold higher than at present and lows at approximately 200 ppm (60-62).

Ice-core records are reported to show seven extended periods during 650,000 years in which CO₂, methane (CH₄), and temperature increased and then decreased (63-65). Ice-core records contain substantial uncertainties (58), so these correlations are imprecise.

In all seven glacial and interglacial cycles, the reported changes in CO₂ and CH₄ lagged the temperature changes and could not, therefore, have caused them (66). These fluctuations probably involved temperature-caused changes in oceanic and terrestrial CO₂ and CH₄ content. More recent CO₂ fluctuations also lag temperature (67,68).

In 1957, Revelle and Seuss (69) estimated that temperature-caused out-gassing of ocean CO₂ would increase atmospheric




Figure 16: Temperature rise versus CO₂ rise from seven ice-core measured interglacial periods (63-65); from calculations (69) and measurements (70) of sea water out-gassing; and as measured during the 20th and 21st centuries (10,72). The interglacial temperature increases caused the CO₂ rises through release of ocean CO₂. The CO₂ rises did not cause the temperature rises.

In addition to the agreement between the out-gassing estimates and measurements, this conclusion is also verified by the small temperature rise during the 20th and 21st centuries. If the CO₂ versus temperature correlation during the seven interglacials had been caused by CO₂ greenhouse warming, then the temperature rise per CO₂ rise would have been as high during the 20th and 21st centuries as it was during the seven interglacial periods.




Figure 17: Atmospheric CO₂ concentrations in parts per million by volume, ppm, measured spectrophotometrically at Mauna Loa, Hawaii, between 1958 and 2007. These measurements agree well with those at other locations (71). Data before 1958 are from ice cores and chemical analyses, which have substantial experimental uncertainties. We have used 295 ppm for the period 1880 to 1890, which is an average of the available estimates. About 0.6 Gt C of CO₂ is produced annually by human respiration and often leads to concentrations exceeding 1,000 ppm in public buildings. Atmospheric CO₂ has increased 22% since 1958 and about 30% since 1880.

CO₂ by about 7% per °C temperature rise. The reported change during the seven interglacials of the 650,000-year ice core record is about 5% per °C (63), which agrees with the out-gassing calculation.

Between 1900 and 2006, Antarctic CO₂ increased 30% per 0.1 °C temperature change (72), and world CO₂ increased 30% per 0.5 °C. In addition to ocean out-gassing, CO₂ from human use of hydrocarbons is a new source. Neither this new source nor the older natural CO₂ sources are causing atmospheric temperature to change.

The hypothesis that the CO₂ rise during the interglacials caused the temperature to rise requires an increase of about 6 °C per 30% rise in CO₂ as seen in the ice core record. If this hypothesis were correct, Earth temperatures would have risen about 6 °C between 1900 and 2006, rather than the rise of between 0.1 °C and 0.5 °C, which actually occurred. This difference is illustrated in Figure 16.

The 650,000-year ice-core record does not, therefore, agree with the hypothesis of "human-caused global warming," and, in fact, provides empirical evidence that invalidates this hypothesis.

Carbon dioxide has a very short residence time in the atmosphere. Beginning with the 7 to 10-year half-time of CO₂ in the atmosphere estimated by Revelle and Seuss (69), there were 36 estimates of the atmospheric CO₂ half-time based upon experimental measurements published between 1957 and 1992 (59). These range between 2 and 25 years, with a mean of 7.5, a median of 7.6, and an upper range average of about 10. Of the 36 values, 33 are 10 years or less.

Many of these estimates are from the decrease in atmospheric carbon 14 after cessation of atmospheric nuclear weapons testing, which provides a reliable half-time. There is no experimental evidence to support computer model estimates (73) of a CO₂ atmospheric "lifetime" of 300 years or more.

Human production of 8 Gt C per year of CO₂ is negligible as compared with the 40,000 Gt C residing in the oceans and biosphere. At ultimate equilibrium, human-produced CO₂ will have an insignificant effect on the amounts in the various reservoirs. The rates of approach to equilibrium are, however, slow enough that human use creates a transient atmospheric increase.

In any case, the sources and amounts of CO₂ in the atmosphere are of secondary importance to the hypothesis of "human-caused global warming." It is human burning of coal, oil, and natural gas that is at issue. CO₂ is merely an intermediate in a hypothetical mechanism by which this "human-caused global warming" is said to take place. The amount of atmospheric CO₂ does have profound environmental effects on plant and animal populations (74) and diversity, as is discussed below.

CLIMATE CHANGE

While the average temperature change taking place as the Earth recovers from the Little Ice Age is so slight that it is difficult to discern, its environmental effects are measurable. Glacier shortening and the 7 inches per century rise in sea level are examples. There are additional climate changes that are correlated with this rise in temperature and may be caused by it.

Greenland, for example, is beginning to turn green again, as it was 1,000 years ago during the Medieval Climate Optimum (11). Arctic sea ice is decreasing somewhat (75), but Antarctic ice is not decreasing and may be increasing, due to increased snow (76-79).

In the United States, rainfall is increasing at about 1.8 inches per century, and the number of severe tornados is decreasing, as shown in Figures 7 and 8. If world temperatures continue to rise at the current rate, they will reach those of the Medieval Climate Optimum about 2 centuries from now. Historical reports of that period record the growing of warm weather crops in localities too cold for that purpose today, so it is to be expected that the area of more temperate climate will expand as it did then. This is already being observed, as studies at higher altitudes have reported increases in amount and diversity of plant and animal life by more than 50% (12,80).

Atmospheric temperature is increasing more in the Northern Hemisphere than in the Southern, with intermediate periods of increase and decrease in the overall trends.

There has been no increase in frequency or severity of Atlantic hurricanes during the period of 6-fold increase in hydrocarbon use, as is illustrated in Figures 9 and 10. Numbers of violent hurricanes vary greatly from year to year and are no greater now than they were 50 years ago. Similarly, maximum wind speeds have not increased.

All of the observed climate changes are gradual, moderate, and entirely within the bounds of ordinary natural changes that have occurred during the benign period of the past few thousand years.

There is no indication whatever in the experimental data that an abrupt or remarkable change in any of the ordinary natural climate variables is beginning or will begin to take place.

GLOBAL WARMING HYPOTHESIS

The greenhouse effect amplifies solar warming of the earth. Greenhouse gases such as H₂O, CO₂, and CH₄ in the Earth's atmosphere, through combined convective readjustments and the radiative blanketing effect, essentially decrease the net escape of terrestrial thermal infrared radiation. Increasing CO₂, therefore, effectively increases radiative energy input to the Earth's atmosphere. The path of this radiative input is complex. It is redistributed, both vertically and horizontally, by various physical processes, including advection, convection, and diffusion in the atmosphere and ocean.

When an increase in CO₂ increases the radiative input to the atmosphere, how and in which direction does the atmosphere respond? Hypotheses about this response differ and are schematically shown in Figure 18. Without the water-vapor greenhouse effect, the Earth would be about 14 °C cooler (81). The radiative contribution of doubling atmospheric CO₂ is minor, but this radiative greenhouse effect is treated quite differently by different climate hypotheses. The hypotheses that the IPCC (82,83) has chosen to adopt predict that the effect of CO₂ is amplified by the atmosphere, especially by water vapor, to produce a large temperature increase. Other hypotheses, shown as hypothesis 2, predict the opposite – that the atmospheric response will counteract the CO₂ increase and result in insignificant changes in global temperature (81,84,85,91,92). The experimental evidence, as described above, favors hypothesis 2. While CO₂ has increased substantially, its effect on temperature has been so slight that it has not been experimentally detected.

The computer climate models upon which "human-caused global warming" is based have substantial uncertainties and are markedly unreliable. This is not surprising, since the climate is a coupled,




Figure 18: Qualitative illustration of greenhouse warming. "Present GHE" is the current greenhouse effect from all atmospheric phenomena. "Radiative effect of CO₂" is the added greenhouse radiative effect from doubling CO₂ without consideration of other atmospheric components. "Hypothesis 1 IPCC" is the hypothetical amplification effect assumed by IPCC. "Hypothesis 2" is the hypothetical moderation effect.

non-linear dynamical system. It is very complex. Figure 19 illustrates the difficulties by comparing the radiative CO₂ greenhouse effect with correction factors and uncertainties in some of the parameters in the computer climate calculations. Other factors, too, such as the chemical and climatic influence of volcanoes, cannot now be reliably computer modeled.

In effect, an experiment has been performed on the Earth during the past half-century – an experiment that includes all of the complex factors and feedback effects that determine the Earth's temperature and climate. Since 1940, hydrocarbon use has risen 6-fold. Yet, this rise has had no effect on the temperature trends, which have continued their cycle of recovery from the Little Ice Age in close correlation with increasing solar activity.

Not only has the global warming hypothesis failed experimental tests, it is theoretically flawed as well. It can reasonably be argued that cooling from negative physical and biological feedbacks to greenhouse gases nullifies the slight initial temperature rise (84,86).

The reasons for this failure of the computer climate models are subjects of scientific debate (87). For example, water vapor is the largest contributor to the overall greenhouse effect (88). It has been suggested that the climate models treat feedbacks from clouds, water vapor, and related hydrology incorrectly (85,89-92).

The global warming hypothesis with respect to CO₂ is not based upon the radiative properties of CO₂ itself, which is a very weak greenhouse gas. It is based upon a small initial increase in temperature caused by CO₂ and a large theoretical amplification of that temperature increase, primarily through increased evaporation of H₂O, a




Figure 19: The radiative greenhouse effect of doubling the concentration of atmospheric CO₂ (right bar) as compared with four of the uncertainties in the computer climate models (87,93).




Figure 20: Global atmospheric methane concentration in parts per million between 1982 and 2004 (94).

strong greenhouse gas. Any comparable temperature increase from another cause would produce the same calculated outcome.

Thus, the 3,000-year temperature record illustrated in Figure 1 also provides a test of the computer models. The historical temperature record shows that the Earth has previously warmed far more than could be caused by CO₂ itself. Since these past warming cycles have not initiated water-vapor-mediated atmospheric warming catastrophes, it is evident that weaker effects from CO₂ cannot do so.

Methane is also a minor greenhouse gas. World CH₄ levels are, as shown in Figure 20, leveling off. In the U.S. in 2005, 42% of human-produced methane was from hydrocarbon energy production, 28% from waste management, and 30% from agriculture (95). The total amount of CH₄ produced from these U.S. sources decreased 7% between 1980 and 2005. Moreover, the record shows that, even while methane was increasing, temperature trends were benign.

The “human-caused global warming” – often called the “global warming” – hypothesis depends entirely upon computer model-generated scenarios of the future. There are no empirical records that verify either these models or their flawed predictions (96).

Claims (97) of an epidemic of insect-borne diseases, extensive species extinction, catastrophic flooding of Pacific islands, ocean acidification, increased numbers and severities of hurricanes and tornados, and increased human heat deaths from the 0.5 °C per century temperature rise are not consistent with actual observations. The “human-caused global warming” hypothesis and the computer calculations that support it are in error. They have no empirical support and are invalidated by numerous observations.

WORLD TEMPERATURE CONTROL

World temperature is controlled by natural phenomena. What steps could mankind take if solar activity or other effects began to shift the Earth toward temperatures too cold or too warm for optimum human life?

First, it would be necessary to determine what temperature humans feel is optimum. It is unlikely that the chosen temperature would be exactly that which we have today. Second, we would be fortunate if natural forces were to make the Earth too warm rather than too cold because we can cool the Earth with relative ease. We have no means by which to warm it. Attempting to warm the Earth with addition of CO₂ or to cool the Earth by restrictions of CO₂ and hydrocarbon use would, however, be futile. Neither would work.

Inexpensively blocking the sun by means of particles in the upper atmosphere would be effective. S.S. Penner, A.M. Schneider, and E. M. Kennedy have proposed (98) that the exhaust systems of commercial airliners could be tuned in such a way as to eject particulate sun-blocking material into the upper atmosphere. Later, Edward Teller similarly suggested (18) that particles could be injected into

the atmosphere in order to reduce solar heating and cool the Earth. Teller estimated a cost of between \$500 million and \$1 billion per year for between 1 °C and 3 °C of cooling. Both methods use particles so small that they would be invisible from the Earth.

These methods would be effective and economical in blocking solar radiation and reducing atmospheric and surface temperatures. There are other similar proposals (99). World energy rationing, on the other hand, would not work.

The climate of the Earth is now benign. If temperatures become too warm, this can easily be corrected. If they become too cold, we have no means of response – except to maximize nuclear and hydrocarbon energy production and technological advance. This would help humanity adapt and might lead to new mitigation technology.

FERTILIZATION OF PLANTS BY CO₂

How high will the CO₂ concentration of the atmosphere ultimately rise if mankind continues to increase the use of coal, oil, and natural gas? At ultimate equilibrium with the ocean and other reservoirs there will probably be very little increase. The current rise is a non-equilibrium result of the rate of approach to equilibrium.

One reservoir that would moderate the increase is especially important. Plant life provides a large sink for CO₂. Using current knowledge about the increased growth rates of plants and assuming increased CO₂ release as compared to current emissions, it has been estimated that atmospheric CO₂ levels may rise to about 600 ppm before leveling off. At that level, CO₂ absorption by increased Earth biomass is able to absorb about 10 Gt C per year (100). At present, this absorption is estimated to be about 3 Gt C per year (57).

About 30% of this projected rise from 295 to 600 ppm has already taken place, without causing unfavorable climate changes. Moreover, the radiative effects of CO₂ are logarithmic (101,102), so more than 40% of any climatic influences have already occurred.

As atmospheric CO₂ increases, plant growth rates increase. Also, leaves transpire less and lose less water as CO₂ increases, so that plants are able to grow under drier conditions. Animal life, which depends upon plant life for food, increases proportionally.

Figures 21 to 24 show examples of experimentally measured increases in the growth of plants. These examples are representative of a very large research literature on this subject (103-109). As Figure 21 shows, long-lived 1,000- to 2,000-year-old pine trees have shown a sharp increase in growth during the past half-century. Figure 22 shows the 40% increase in the forests of the United States that has




Figure 21: Standard deviation from the mean of tree ring widths for (a) bristlecone pine, limber pine, and fox tail pine in the Great Basin of California, Nevada, and Arizona and (b) bristlecone pine in Colorado (110). Tree ring widths were averaged in 20-year segments and then normalized so that the means of prior tree growth were zero. The deviations from the means are shown in units of standard deviations of those means.




Figure 22: Inventories of standing hardwood and softwood timber in the United States compiled in *Forest Resources of the United States, 2002*, U.S. Department of Agriculture Forest Service (111,112). The linear trend cited in 1998 (1) with an increase of 30% has continued. The increase is now 40%. The amount of U.S. timber is rising almost 1% per year.

taken place since 1950. Much of this increase is due to the increase in atmospheric CO₂ that has already occurred. In addition, it has been reported that Amazonian rain forests are increasing their vegetation by about 900 pounds of carbon per acre per year (113), or approximately 2 tons of biomass per acre per year. Trees respond to CO₂ fertilization more strongly than do most other plants, but all plants respond to some extent.

Since plant response to CO₂ fertilization is nearly linear with respect to CO₂ concentration over the range from 300 to 600 ppm, as seen in Figure 23, experimental measurements at different levels of CO₂ enrichment can be extrapolated. This has been done in Figure 24 in order to illustrate CO₂ growth enhancements calculated for the atmospheric increase of about 88 ppm that has already taken place and those expected from a projected total increase of 305 ppm.

Wheat growth is accelerated by increased atmospheric CO₂, especially under dry conditions. Figure 24 shows the response of wheat grown under wet conditions versus that of wheat stressed by lack of water. The underlying data is from open-field experiments. Wheat was grown in the usual way, but the atmospheric CO₂ concentrations of circular sections of the fields were increased by arrays of com-




Figure 23: Summary data from 279 published experiments in which plants of all types were grown under paired stressed (open red circles) and unstressed (closed blue circles) conditions (114). There were 208, 50, and 21 sets at 300, 600, and an average of about 1350 ppm CO₂, respectively. The plant mixture in the 279 studies was slightly biased toward plant types that respond less to CO₂ fertilization than does the actual global mixture. Therefore, the figure underestimates the expected global response. CO₂ enrichment also allows plants to grow in drier regions, further increasing the response.

puter-controlled equipment that released CO₂ into the air to hold the levels as specified (115,116). Orange and young pine tree growth enhancement (117-119) with two atmospheric CO₂ increases – that which has already occurred since 1885 and that projected for the next two centuries – is also shown. The relative growth enhancement of trees by CO₂ diminishes with age. Figure 24 shows young trees.

Figure 23 summarizes 279 experiments in which plants of various types were raised under CO₂-enhanced conditions. Plants under stress from less-than-ideal conditions – a common occurrence in nature – respond more to CO₂ fertilization. The selections of species in Figure 23 were biased toward plants that respond less to CO₂ fertilization than does the mixture actually covering the Earth, so Figure 23 underestimates the effects of global CO₂ enhancement.

Clearly, the green revolution in agriculture has already benefitted from CO₂ fertilization, and benefits in the future will be even greater. Animal life is increasing proportionally, as shown by studies of 51 terrestrial (120) and 22 aquatic ecosystems (121). Moreover, as shown by a study of 94 terrestrial ecosystems on all continents ex-




Figure 24: Calculated (1,2) growth rate enhancement of wheat, young orange trees, and very young pine trees already taking place as a result of atmospheric enrichment by CO₂ from 1885 to 2007 (a), and expected as a result of atmospheric enrichment by CO₂ to a level of 600 ppm (b).

cept Antarctica (122), species richness – biodiversity – is more positively correlated with productivity – the total quantity of plant life per acre – than with anything else.

Atmospheric CO₂ is required for life by both plants and animals. It is the sole source of carbon in all of the protein, carbohydrate, fat, and other organic molecules of which living things are constructed.

Plants extract carbon from atmospheric CO₂ and are thereby fertilized. Animals obtain their carbon from plants. Without atmospheric CO₂, none of the life we see on Earth would exist.

Water, oxygen, and carbon dioxide are the three most important substances that make life possible.

They are surely not environmental pollutants.

ENVIRONMENT AND ENERGY

The single most important human component in the preservation of the Earth's environment is energy. Industrial conversion of energy into forms that are useful for human activities is the most important aspect of technology. Abundant inexpensive energy is required for the prosperous maintenance of human life and the continued advance of life-enriching technology. People who are prosperous have the wealth required to protect and enhance their natural environment.

Currently, the United States is a net importer of energy as shown in Figure 25. Americans spend about \$300 billion per year for imported oil and gas – and an additional amount for military expenses related to those imports.




Figure 25: In 2006, the United States obtained 84.9% of its energy from hydrocarbons, 8.2% from nuclear fuels, 2.9% from hydroelectric dams, 2.1% from wood, 0.8% from biofuels, 0.4% from waste, 0.3% from geothermal, and 0.3% from wind and solar radiation. The U.S. uses 21 million barrels of oil per day – 27% from OPEC, 17% from Canada and Mexico, 16% from others, and 40% produced in the U.S. (95). The cost of imported oil and gas at \$60 per barrel and \$7 per 1,000 ft³ in 2007 is about \$300 billion per year.

Political calls for a reduction of U.S. hydrocarbon use by 90% (123), thereby eliminating 75% of America's energy supply, are obviously impractical. Nor can this 75% of U.S. energy be replaced by alternative "green" sources. Despite enormous tax subsidies over the past 30 years, green sources still provide only 0.3% of U.S. energy.

Yet, the U.S. clearly cannot continue to be a large net importer of energy without losing its economic and industrial strength and its political independence. It should, instead, be a net exporter of energy.

There are three realistic technological paths to American energy independence – increased use of hydrocarbon energy, nuclear energy, or both. There are no climatological impediments to increased use of hydrocarbons, although local environmental effects can and must be accommodated. Nuclear energy is, in fact, less expensive and more environmentally benign than hydrocarbon energy, but it too has been the victim of the politics of fear and claimed disadvantages and dangers that are actually negligible.

For example, the "problem" of high-level "nuclear waste" has been given much attention, but this problem has been politically created by U.S. government barriers to American fuel breeding and reprocessing. Spent nuclear fuel can be recycled into new nuclear fuel. It need not be stored in expensive repositories.

Reactor accidents are also much publicized, but there has never been even one human death associated with an American nuclear reactor incident. By contrast, American dependence on automobiles results in more than 40,000 human deaths per year.

All forms of energy generation, including "green" methods, entail industrial deaths in the mining, manufacture, and transport of resources they require. Nuclear energy requires the smallest amount of such resources (124) and therefore has the lowest risk of deaths.

Estimated relative costs of electrical energy production vary with

geographical location and underlying assumptions. Figure 26 shows a recent British study, which is typical. At present, 43% of U.S. energy consumption is used for electricity production.

To be sure, future inventions in energy technology may alter the relative economics of nuclear, hydrocarbon, solar, wind, and other methods of energy generation. These inventions cannot, however, be forced by political fiat, nor can they be wished into existence. Alternatively, "conservation," if practiced so extensively as to be an alternative to hydrocarbon and nuclear power, is merely a politically correct word for "poverty."

The current untenable situation in which the United States is losing \$300 billion per year to pay for foreign oil and gas is not the result of failures of government energy production efforts. The U.S. government does not produce energy. Energy is produced by private industry. Why then has energy production thrived abroad while domestic production has stagnated?

This stagnation has been caused by United States government taxation, regulation, and sponsorship of litigation, which has made the U.S. a very unfavorable place to produce energy. In addition, the U.S. government has spent vast sums of tax money subsidizing inferior energy technologies for political purposes.

It is not necessary to discern in advance the best course to follow. Legislative repeal of taxation, regulation, incentives to litigation, and repeal of all subsidies of energy generation industries would stimulate industrial development, wherein competition could then automatically determine the best paths.

Nuclear power is safer, less expensive, and more environmentally benign than hydrocarbon power, so it is probably the better choice for increased energy production. Solid, liquid and gaseous hydrocarbon fuels provide, however, many conveniences, and a national infrastructure to use them is already in place. Oil from shale or coal liquefaction is less expensive than crude oil at current prices, but its ongoing production costs are higher than those for already developed oil fields. There is, therefore, an investment risk that crude oil prices could drop so low that liquefaction plants could not compete. Nuclear energy does not have this disadvantage, since the operating costs of nuclear power plants are very low.

Figure 27 illustrates, as an example, one practical and environmentally sound path to U.S. energy independence. At present 19% of U.S. electricity is produced by 104 nuclear power reactors with an average generating output in 2006 of 870 megawatts per reactor, for a total of about 90 GWe (gigawatts) (125). If this were increased by 560 GWe, nuclear power could fill all current U.S. electricity requirements and have 230 GWe left over for export as electricity or as hydrocarbon fuels replaced or manufactured.

Thus, rather than a \$300 billion trade loss, the U.S. would have a \$200 billion trade surplus – and installed capacity for future U.S. re-




Figure 26: Delivered cost per kilowatt hour of electrical energy in Great Britain in 2006, without CO₂ controls (126). These estimates include all capital and operational expenses for a period of 50 years. Micro wind or solar are units installed for individual homes.




Figure 27: Construction of one Palo Verde installation with 10 reactors in each of the 50 states. Energy trade deficit is reversed by \$500 billion per year, resulting in a \$200 billion annual surplus. Currently, this solution is not possible owing to misguided government policies, regulations, and taxation and to legal maneuvers available to anti-nuclear activists. These impediments should be legislatively repealed.

quirements. Moreover, if heat from additional nuclear reactors were used for coal liquefaction and gasification, the U.S. would not even need to use its oil resources. The U.S. has about 25% of the world's coal reserves. This heat could also liquify biomass, trash, or other sources of hydrocarbons that might eventually prove practical.

The Palo Verde nuclear power station near Phoenix, Arizona, was originally intended to have 10 nuclear reactors with a generating capacity of 1,243 megawatts each. As a result of public hysteria caused by false information – very similar to the human-caused global warming hysteria being spread today, construction at Palo Verde was stopped with only three operating reactors completed. This installation is sited on 4,000 acres of land and is cooled by waste water from the city of Phoenix, which is a few miles away. An area of 4,000 acres is 6.25 square miles or 2.5 miles square. The power station itself occupies only a small part of this total area.

If just one station like Palo Verde were built in each of the 50 states and each installation included 10 reactors as originally planned for Palo Verde, these plants, operating at the current 90% of design capacity, would produce 560 GWe of electricity. Nuclear technology has advanced substantially since Palo Verde was built, so plants constructed today would be even more reliable and efficient.

Assuming a construction cost of \$2.3 billion per 1,200 MWe reactor (127) and 15% economies of scale, the total cost of this entire project would be \$1 trillion, or 4 months of the current U.S. federal budget. This is 8% of the annual U.S. gross domestic product. Construction costs could be repaid in just a few years by the capital now spent by the people of the United States for foreign oil and by the change from U.S. import to export of energy.

The 50 nuclear installations might be sited on a population basis. If so, California would have six, while Oregon and Idaho together would have one. In view of the great economic value of these facilities, there would be vigorous competition for them.

In addition to these power plants, the U.S. should build fuel reprocessing capability, so that spent nuclear fuel can be reused. This would lower fuel cost and eliminate the storage of high-level nuclear waste. Fuel for the reactors can be assured for 1,000 years (128) by using both ordinary reactors with high breeding ratios and specific breeder reactors, so that more fuel is produced than consumed.

About 33% of the thermal energy in an ordinary nuclear reactor is converted to electricity. Some new designs are as high as 48%. The heat from a 1,243 MWe reactor can produce 38,000 barrels of coal-derived oil per day (129). With one additional Palo Verde installation in each state for oil production, the yearly output would be at least 7 billion barrels per year with a value, at \$60 per barrel, of

more than \$400 billion per year. This is twice the oil production of Saudi Arabia. Current proven coal reserves of the United States are sufficient to sustain this production for 200 years (128). This liquified coal exceeds the proven oil reserves of the entire world. The reactors could produce gaseous hydrocarbons from coal, too.

The remaining heat from nuclear power plants could warm air or water for use in indoor climate control and other purposes.

Nuclear reactors can also be used to produce hydrogen, instead of oil and gas (130,131). The current cost of production and infrastructure is, however, much higher for hydrogen than for oil and gas. Technological advance reduces cost, but usually not abruptly. A prescient call in 1800 for the world to change from wood to methane would have been impracticably ahead of its time, as may be a call today for an abrupt change from oil and gas to hydrogen. In distinguishing the practical from the futuristic, a free market in energy is absolutely essential.

Surely these are better outcomes than are available through international rationing and taxation of energy as has been recently proposed (82,83,97,123). This nuclear energy example demonstrates that current technology can produce abundant inexpensive energy if it is not politically suppressed.

There need be no vast government program to achieve this goal. It could be reached simply by legislatively removing all taxation, most regulation and litigation, and all subsidies from all forms of energy production in the U.S., thereby allowing the free market to build the most practical mixture of methods of energy generation.

With abundant and inexpensive energy, American industry could be revitalized, and the capital and energy required for further industrial and technological advance could be assured. Also assured would be the continued and increased prosperity of all Americans.

The people of the United States need more low-cost energy, not less. If this energy is produced in the United States, it can not only become a very valuable export, but it can also ensure that American industry remains competitive in world markets and that hoped-for American prosperity continues and grows.

In this hope, Americans are not alone. Across the globe, billions of people in poorer nations are struggling to improve their lives. These people need abundant low-cost energy, which is the currency of technological progress.

In newly developing countries, that energy must come largely from the less technologically complicated hydrocarbon sources. It is a moral imperative that this energy be available. Otherwise, the efforts of these peoples will be in vain, and they will slip backwards into lives of poverty, suffering, and early death.


Energy is the foundation of wealth. Inexpensive energy allows people to do wonderful things. For example, there is concern that it may become difficult to grow sufficient food on the available land. Crops grow more abundantly in a warmer, higher CO₂ environment, so this can mitigate future problems that may arise (12).

Energy provides, however, an even better food insurance plan. Energy-intensive hydroponic greenhouses are 2,000 times more productive per unit land area than are modern American farming methods (132). Therefore, if energy is abundant and inexpensive, there is no practical limit to world food production.

Fresh water is also believed to be in short supply. With plentiful inexpensive energy, sea water desalination can provide essentially unlimited supplies of fresh water.

During the past 200 years, human ingenuity in the use of energy has produced many technological miracles. These advances have markedly increased the quality, quantity, and length of human life. Technologists of the 21st century need abundant, inexpensive energy with which to continue this advance.

Were this bright future to be prevented by world energy rationing, the result would be tragic indeed. In addition to human loss, the Earth's environment would be a major victim of such a mistake. Inexpensive energy is essential to environmental health. Prosperous people have the wealth to spare for environmental preservation and enhancement. Poor, impoverished people do not.



Global monthly average lower troposphere temperature since 1979 according to [University of Alabama](#) at Huntsville (UAH), USA. This graph uses data obtained by the National Oceanographic and Atmospheric Administration (NOAA) TIROS-N satellite, interpreted by [Dr. Roy Spencer](#) and [Dr. John Christy](#), both at Global Hydrology and Climate Center, [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37 month average, nearly corresponding to a running 3 yr average. The cooling and warming periods directly influenced by [the 1991 Mt. Pinatubo volcanic eruption](#) and the 1998 El Niño, respectively, are clearly visible. Reference period 1981-2010. Last month shown: December 2016. Last diagram update: 4 January 2017.

- [Click here](#) to see the latest UAH MSU global monthly lower troposphere temperature anomaly with comments.
- [Click here](#) to download the series of UAH MSU global monthly lower troposphere temperature anomalies since December 1978.
- [Click here](#) to see a maturity diagram for the MSU UAH data series.
- [Click here](#) to read about data smoothing.
- [Click here](#) to read about the latest version (6.0) of the UAH Temperature Dataset (April 28, 2015).


Schedule DWS - 3

Start Download - View PDF

Convert From Doc to PDF, PDF to Doc Simply With The Free On-line App! Go to fromdoctopdf.com



Latest Global Temps



Latest Global Average Tropospheric Temperatures

Since 1979, NOAA satellites have been carrying instruments which measure the natural microwave thermal emissions from oxygen in the atmosphere. The intensity of the signals these microwave radiometers measure at different microwave frequencies is directly proportional to the temperature of different, deep layers of the atmosphere. Every month, John Christy and I update global temperature datasets that represent the piecing together of the temperature data from a total of fourteen instruments flying on different satellites over the years. A discussion of the latest version (6.0) of the dataset is located [here](#).

The graph above represents the latest update; updates are usually made within the first week of every month. Contrary to some reports, the satellite measurements are not calibrated in any way with the global surface-based thermometer records of temperature. They instead use their own on-board precision redundant platinum resistance thermometers (PRTs) calibrated to a laboratory reference standard before launch.

© Roy Spencer, PhD